

1107
PRV

PATENT- OCH REGISTRERINGSVERKET
Patentavdelningen

09/647 544

PCT/SE 99/00544


**Intyg
Certificate**



Härmed intygas att bifogade kopior överensstämmer med de handlingar som ursprungligen ingivits till Patent- och registreringsverket i nedannämnda ansökan.

This is to certify that the annexed is a true copy of the documents as originally filed with the Patent- and Registration Office in connection with the following patent application.

(71) Sökande *Sten Trolle, Lund, SE*
Applicant (s)



(21) Patentansökningsnummer *9801164-6*
Patent application number

(86) Ingivningsdatum *1998-04-02*
Date of filing

Stockholm, 1999-07-05

**PRIORITY
DOCUMENT**

SUBMITTED OR TRANSMITTED IN
COMPLIANCE WITH RULE 17.1(a) OR (b)

*För Patent- och registreringsverket
For the Patent- and Registration Office*

Evy Morin
Evy Morin

Avgift
Fee

ISOLATION, CLONING AND SEQUENCE ANALYSIS OF THE INTEGRIN
SUBUNIT $\alpha 10$, A $\beta 1$ -ASSOCIATED COLLAGEN-BINDING INTEGRIN
EXPRESSED ON CHONDROCYTES

Lisbet Camper¹, Ulf Hellman² and Evy Lundgren-Åkerlund¹

¹Department of Cell and Molecular Biology, Section for Cell and Matrix Biology, Lund
University, Lund Sweden

²Ludwig Institute for Cancer Research, Uppsala, Sweden

Address all correspondence to:

Lisbet Camper

Department of Cell and Molecular Biology, Section for Cell and Matrix Biology

Lund University, P.O. Box 94, S-221 00 Lund, Sweden

Telephone: +46 (46) 222 3126; FAX: +46 (46) 222 3128

e-mail: lisbet.camper@medkem.lu.se

Running title: Integrin $\alpha 10\beta 1$

Key word: integrin, $\alpha 10\beta 1$, chondrocytes, collagen, sequence analysis

Abstract

We have found that chondrocytes express a novel collagen type II-binding integrin, a new member of the $\beta 1$ -integrin family. The integrin α -subunit, which has a Mr of 160 kD (reducing conditions), was isolated from bovine chondrocytes by affinity purification. The human homologue was obtained by screening a human chondrocyte library with a bovine cDNA probe. Cloning and cDNA sequence analysis of the human integrin α -subunit, designated $\alpha 10$, show that it shares the general structure of other integrin α -subunits. The predicted amino acid sequence consists of a 1167 amino acid mature protein, including a signal peptide (22 aa), a long extracellular domain (1098 aa), a transmembrane domain (25 aa) and a short cytoplasmic domain (22 aa). The extracellular part contains a seven-fold repeated sequence, an I-domain (199 aa) and three putative divalent cation-binding sites. The deduced amino acid sequence of $\alpha 10$ is 35% identical to the integrin subunit $\alpha 2$ and 37% identical to the integrin subunit $\alpha 1$. In northern blot analysis a single mRNA of 5.4 kb was detected in chondrocytes. A peptide antibody against the predicted sequence of the cytoplasmic domain of $\alpha 10$, immunoprecipitated two proteins with M_s of 125 kD and 160 kD from chondrocyte lysates under reducing conditions. The peptide antibody specifically stained chondrocytes in tissue sections of human articular cartilage showing that $\alpha 10\beta 1$ indeed is expressed in cartilage tissue.

Introduction

The integrins are a large family of transmembrane glycoproteins that mediate cell-cell and cell-matrix interactions (1-5). All known members of this superfamily are non-covalently associated heterodimers composed of an α - and a β - subunit. At present, 8 β - ($\beta 1$ - $\beta 8$) (for references see (6)) and 16 α - subunits ($\alpha 1$ - $\alpha 9$, αv , αM , αL , αX , αIIb , αE and αD) have been characterised (6-21) and these subunits associate to generate more than 20 different integrins. The $\beta 1$ -subunit has been shown to associate with ten different α -subunits, $\alpha 1$ - $\alpha 9$ and αv , and to mediate interactions with extracellular matrix proteins such as collagens, laminins and fibronectin. The major collagen binding integrins are $\alpha 1\beta 1$ and $\alpha 2\beta 1$ (22-25). The integrins $\alpha 3\beta 1$ and $\alpha 9\beta 1$ have also been reported to interact with collagen (26;27) although this interaction is not well understood (28). The extracellular N-terminal regions of the α and β integrin subunits are important in the binding of ligands (29;30). The N-terminal region of the α -subunits is composed of a seven-fold repeated sequence (12;31) containing FG and GAP consensus sequences. The repeats are predicted to fold into a β -propeller domain (32) with the last three or four repeats containing putative divalent cation binding sites. The α -integrin subunits $\alpha 1$, $\alpha 2$, αD , αE , αL , αM and αX contain a ~200 amino acid inserted domain, the I-domain (A-domain), which shows similarity to sequences in von Willebrand factor, cartilage matrix protein and complement factors C2 and B (33;34). The I-domain is localised between the second and third FG-GAP repeats, it contains a metal ion-dependent adhesion site (MIDAS) and it is involved in binding of ligands (35-38).

Chondrocytes, the only type of cells in cartilage, express a number of different integrins including $\alpha 1\beta 1$, $\alpha 2\beta 1$, $\alpha 3\beta 1$, $\alpha 5\beta 1$, $\alpha 6\beta 1$, $\alpha v\beta 3$, and $\alpha v\beta 5$ (39-41). We have shown that $\alpha 1\beta 1$ and $\alpha 2\beta 1$ mediate chondrocyte interactions with collagen type II (25) which is one of the major components in cartilage. We have also shown that $\alpha 2\beta 1$ is a receptor for the

cartilage matrix protein chondroadherin (42). In the present study we have isolated a novel collagen type II binding integrin, $\alpha 10\beta 1$, from bovine articular chondrocytes. Cloning and sequence analysis of the human homologue is described and expression of $\alpha 10$ on chondrocytes is examined.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038
1039
1040
1041
1042
1043
1044
1045
1046
1047
1048
1049
1050
1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1120
1121
1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180
1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197
1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285
1286
1287
1288
1289
1290
1291
1292
1293
1294
1295
1296
1297
1298
1299
1300
1301
1302
1303
1304
1305
1306
1307
1308
1309
1310
1311
1312
1313
1314
1315
1316
1317
1318
1319
1320
1321
1322
1323
1324
1325
1326
1327
1328
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1340
1341
1342
1343
1344
1345
1346
1347
1348
1349
1350
1351
1352
1353
1354
1355
1356
1357
1358
1359
1360
1361
1362
1363
1364
1365
1366
1367
1368
1369
1370
1371
1372
1373
1374
1375
1376
1377
1378
1379
1380
1381
1382
1383
1384
1385
1386
1387
1388
1389
1390
1391
1392
1393
1394
1395
1396
1397
1398
1399
1400
1401
1402
1403
1404
1405
1406
1407
1408
1409
1410
1411
1412
1413
1414
1415
1416
1417
1418
1419
1420
1421
1422
1423
1424
1425
1426
1427
1428
1429
1430
1431
1432
1433
1434
1435
1436
1437
1438
1439
1440
1441
1442
1443
1444
1445
1446
1447
1448
1449
1450
1451
1452
1453
1454
1455
1456
1457
1458
1459
1460
1461
1462
1463
1464
1465
1466
1467
1468
1469
1470
1471
1472
1473
1474
1475
1476
1477
1478
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1530
1531
1532
1533
1534
1535
1536
1537
1538
1539
1540
1541
1542
1543
1544
1545
1546
1547
1548
1549
1550
1551
1552
1553
1554
1555
1556
1557
1558
1559
1560
1561
1562
1563
1564
1565
1566
1567
1568
1569
1570
1571
1572
1573
1574
1575
1576
1577
1578
1579
1580
1581
1582
1583
1584
1585
1586
1587
1588
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1600
1601
1602
1603
1604
1605
1606
1607
1608
1609
1610
1611
1612
1613
1614
1615
1616
1617
1618
1619
1620
1621
1622
1623
1624
1625
1626
1627
1628
1629
1630
1631
1632
1633
1634
1635
1636
1637
1638
1639
1640
1641
1642
1643
1644
1645
1646
1647
1648
1649
1650
1651
1652
1653
1654
1655
1656
1657
1658
1659
1660
1661
1662
1663
1664
1665
1666
1667
1668
1669
1670
1671
1672
1673
1674
1675
1676
1677
1678
1679
1680
1681
1682
1683
1684
1685
1686
1687
1688
1689
1690
1691
1692
1693
1694
1695
1696
1697
1698
1699
1700
1701
1702
1703
1704
1705
1706
1707
1708
1709
1710
1711
1712
1713
1714
1715
1716
1717
1718
1719
1720
1721
1722
1723
1724
1725
1726
1727
1728
1729
1730
1731
1732
1733
1734
1735
1736
1737
1738
1739
1740
1741
1742
1743
1744
1745
1746
1747
1748
1749
1750
1751
1752
1753
1754
1755
1756
1757
1758
1759
1760
1761
1762
1763
1764
1765
1766
1767
1768
1769
1770
1771
1772
1773
1774
1775
1776
1777
1778
1779
1780
1781
1782
1783
1784
1785
1786
1787
1788
1789
1790
1791
1792
1793
1794
1795
1796
1797
1798
1799
1800
1801
1802
1803
1804
1805
1806
1807
1808
1809
1810
1811
1812
1813
1814
1815
1816
1817
1818
1819
1820
1821
1822
1823
1824
1825
1826
1827
1828
1829
1830
1831
1832
1833
1834
1835
1836
1837
1838
1839
1840
1841
1842
1843
1844
1845
1846
1847
1848
1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900
1901
1902
1903
1904
1905
1906
1907
1908
1909
1910
1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058
2059
2060
2061
2062
2063
2064
2065
2066
2067
2068
2069
2070
2071
2072
2073
2074
2075
2076
2077
2078
2079
2080
2081
2082
2083
2084
2085
2086
2087
2088
2089
2090
2091
2092
2093
2094
2095
2096
2097
2098
2099
2100
2101
2102
2103
2104
2105
2106
2107
2108
2109
2110
2111
2112
2113
2114
2115
2116
2117
2118
2119
2120
2121
2122
2123
2124
2125
2126
2127
2128
2129
2130
2131
2132
2133
2134
2135
2136
2137
2138
2139
2140
2141
2142
2143
2144
2145
2146
2147
2148
2149
2150
2151
2152
2153
2154
2155
2156
2157
2158
2159
2160
2161
2162
2163
2164
2165
2166
2167
2168
2169
2170
2171
2172
2173
2174
2175
2176
2177
2178
2179
2180
2181
2182
2183
2184
2185
2186
2187
2188
2189
2190
2191
2192
2193
2194
2195
2196
2197
2198
2199
2200
2201
2202
2203
2204
2205
2206
2207
2208
2209
2210
2211
2212
2213
2214
2215
2216

Materials and Methods

Antibodies

A polyclonal antiserum was generated against the $\alpha 10$ cytoplasmic domain peptide CKKIPEEEKREEKLE. Peptide synthesis, conjugation to KLH, injection of rabbits and affinity purification were performed by Innovagen AB (Lund, Sweden). Monoclonal antibodies against human integrin subunit $\beta 1$ (P4C10), $\alpha 2$ (P1E6) and $\alpha 3$ (P1B5) (unpurified ascites fluid) were from Life Technology Inc, Grand Island NY, USA. The monoclonal antibody against human integrin subunit $\alpha 1$ (TS2/7; hybridoma supernatant) was a kind gift from Timothy Springer, Boston Blood Center, Boston, MA (43). Polyclonal antiserum against the rat $\beta 1$ -integrin subunit was kindly provided by Staffan Johansson, Uppsala, Sweden (44). Polyclonal antibodies (serum) against human integrin subunits $\alpha 2$ (AB1936), $\alpha 3$ (AB1920) and polyclonal antibody (serum) against rat integrin subunits $\alpha 1$ (AB1934) were from Chemicon International Inc. Polyclonal antibodies against the integrin subunit $\alpha 9$ (affinity purified IgG) were a kind gift from Dean Sheppard, UCSF- Lung Biology Center, San Francisco, CA (6).

Cell isolation and culture

Bovine chondrocytes were isolated by digestion of articular cartilage from 4- 6 month old calves with collagenase (CLS1, Worthington) as described elsewhere (45). Briefly, cartilage slices were digested by collagenase in EBSS (Earle's balanced salt solution, GIBCO) for 15-16 hours at 37 °C. The tissue digest was filtered through a 100 μ m nylon filter and the isolated cells were then washed three times in Dulbecco's modified phosphate buffered saline (PBS, GIBCO). Human chondrocytes from articular cartilage were isolated by digestion with pronase (Calbiochem) for 1 hr followed by collagenase (Boehringer Mannheim) for 15-18 hrs, as

described by Häuselmann *et al* (46). The cells were filtered and washed as described above.

Human chondrocytes were cultured in Dulbeccos MEM and F12 (1:1) supplemented with 10%

fetal calf serum, 25 µg/ml ascorbic acid, 50 UI penicillin and 50 µg/ml streptomycin (GIBCO).

To harvest cells, the culture dish was washed three times with Ca/Mg-free PBS and the cells were incubated with 0.5% trypsin and 1 mM EDTA (GIBCO) in PBS (-Ca,Mg) for 5 minutes

Detached cells were suspended in medium containing 10 % FCS or in PBS containing 1mg/ml trypsin inhibitor (Sigma) and then washed in PBS.

Coupling of affinity columns

Collagen type II isolated from nasal cartilage by pepsin digestion (47) was coupled to CNBr-Sepharose (Pharmacia Biotech, Uppsala, Sweden) according to the published procedure (25).

A control column was produced by treating CNBr-Sepharose in a similar manner but in the absence of protein. Bovine fibronectin (Sigma) was coupled to CNBr-Sepharose according to instructions from the manufacturer. After blocking, the fibronectin-Sepharose was washed three times with PBS.

Affinity purification and immunoprecipitation of ¹²⁵I-labeled membrane proteins

Human chondrocyte cell surface proteins were ¹²⁵I-labeled and affinity purified on collagen type II-Sepharose according to the published procedure (25). Cell-lysates or affinity purified samples were immunoprecipitated as described earlier (42). The following antibodies were used in immunoprecipitation experiments: monoclonal antibodies against the human integrin subunits β1, α1, α2 or α3 (unpurified ascites fluid, dilution 1/100); polyclonal antibody against the rat integrin subunit β1 (purified IgG, 50-100 µg/ml); polyclonal peptide antibodies against the integrin subunits α1, α2, α3 and α10 (antiserum, dilution 1/100). The immunoprecipitated

proteins were separated by 4-12% SDS-PAGE and visualized by image analysis using the BioImaging Analyzer Bas2000 (Fuji Photo Film Co., Tokyo Japan).

Affinity purification of the integrin subunit $\alpha 10$ on collagen type II-Sepharose

Freshly isolated bovine chondrocytes (2500×10^6) were lysed in 6 ml of 1 % Triton X-100, 10 mM Tris-HCl, pH 7.4, 100 μ g/ml aprotinin, 2 μ g/ml leupeptin, 2 μ g/ml pepstatin A, 1 mM PMSF, 1 mM $MnCl_2$ and 1 mM $MgCl_2$ for 1 hour on ice. The lysate was centrifuged for 30 minutes at 10,000 rpm and the pellet was discarded. Collagen type II-Sepharose (4 ml) and the fibronectin-Sepharose (2 ml) were equilibrated with at least 20 volumes of 0.1 % Triton X-100, 10 mM Tris-HCl, pH 7.4, 1 mM PMSF, 1 mM $MnCl_2$ and 1 mM $MgCl_2$ (equilibration buffer). The cell-lysate was passed over the fibronectin-Sepharose twice and then incubated with the collagen-Sepharose end over end for 3 hours. The columns were washed (15 gel volumes) with the equilibration buffer containing 75 mM NaCl and bound proteins were eluted (10 x 0.5 ml) with 20 mM EDTA, 1 mM PMSF and 10 mM Tris-HCl, pH 7.4.

Isolation of internal peptides by in-gel digestion and peptide sequencing

Affinity purified proteins were concentrated by precipitation using the methanol/chloroform protocol (48). After reduction/alkylation with DTT/Iodoacetamide (49), samples were subjected to SDS-PAGE on a 4-12 % polyacrylamide gel and protein bands were visualized by Coomassie staining. The 160 kD protein-band was excised from the gel and prepared for in-gel digestion (50). Briefly, the gel slice was washed extensively to remove SDS and the dye, and after complete drying, protease was forced into the gel by rehydration with a solution of either Endoproteinase Lys C (Wako, Neuss, Germany) or modified trypsin (Promega Madison,

WI) in the appropriate digestion buffer. Following an overnight incubation, peptides were extracted and then isolated by narrow bore reversed-phase liquid chromatography on a μ RPC C2/C18 SC 2.1/10 column, operated in a SMART System (Pharmacia Biotech, Uppsala, Sweden). Several peptides were analyzed by Edman degradation in a Perkin-Elmer Applied Biosystem Model 476 sequencer, operated according to the manufacturer's instructions.

mRNA purification and cDNA-synthesis

mRNA from bovine or human chondrocytes were isolated using a QuickPrep Micro mRNA Purification Kit (Pharmacia Biotech, Uppsala, Sweden). cDNA was synthesized at 42° C for 1hr using the Superscript™ II RNase H Reverse Transcriptase cDNA Synthesis system (GIBCO-BRL), random DNA hexamers and oligo (dT) (Promega, Madison, WI).

PCR amplification

PCR-reactions were performed in 50 μ l reaction volumes and contained 1 x Taq polymerase buffer (GIBCO-BRL), 1.5 mM MgCl₂, 1 μ M of each primer, 0.025 U/ μ l Taq polymerase, 1 μ l of DNA template (bovine chondrocyte cDNA) and 0.1 mM each of dATP, dGTP, dCTP and dTTP (Boehringer Mannheim). PCR-samples were heated to 94°C for 5 minutes in a thermocycler and then subjected to 35 cycles consisting of 30 seconds at 94°C (denaturation), 30 seconds at 48°C or 52°C (annealing) and 3 minutes at 72°C (extension). The PCR-products were re-amplified using 1 μ l of each product for an additional 35 cycles. Amplified DNA was analyzed by 1 % agarose gel electrophoresis. Small DNA fragments were analyzed using 4% metha phore agarose (FMC BioProducts).

The degenerate primers GAY AAY ACI GCI CAR AC (DNTAQT, forward) and TIA TIS

WRT GRT GIG GYT (EPHHSI, reverse) were used in PCR to amplify the nucleotide sequence corresponding to the bovine peptide 1 (Table I). A 900 bp PCR-fragment was then amplified from bovine cDNA using an internal specific primer TCA GCC TAC ATT CAG TAT (SAYIQY, forward) corresponding to the cloned nucleotide sequence of peptide 1 together with the degenerate primer ICK RTC CCA RTG ICC IGG (PGHWDR, reverse) corresponding to the bovine peptide 2 (Table I). Mixed bases were used in positions that were twofold degenerate and inosines were used in positions that are three- or fourfold degenerate.

To obtain cDNA that encoded the 5' end of $\alpha 10$ we designed the primer AAC TCG TCT TCC AGT GCC ATT CGT GGG (reverse; residue 1254-1280 in $\alpha 10$ cDNA) and used it for rapid amplification of the cDNA 5' end (RACE) as described in the Marathon[™] cDNA

Amplification kit (Clontech Intermedica).

into bra stor into att det ar humanit.

Cloning and sequencing of cDNA

PCR-fragments were isolated and purified from agarose gels using Jet Sorb DNA extraction kit (Saveen). Purified fragments were then cloned with the Pcr Script[™] Sk(+) kit (Stratagene). Selected plasmides were purified from liquid cultures using QIAGEN-plasmid midi preparation kit (QIAGEN Inc.) and sequenced by ABI 373A- sequencer using ABI Prism[™] Dye Terminator Cycle Sequencing Core kit (PERKIN ELMER) together with T3, T7 and internal specific primers.

Library screening

The cloned 900bp PCR-fragment, corresponding to bovine $\alpha 10$ -integrin, was digoxigenin-labeled according to the DIG DNA labeling kit (Boehringer Mannheim) and used as a probe

for screening of a human articular chondrocyte λ ZapII cDNA library (provided by Dr. Michael Bayliss) (51). Positive clones containing the pBluescript SK⁺ plasmid with the cDNA insert

were rescued from the ZAP vector by *in vivo* excision as described in the ZAP-cDNA® synthesis kit (Stratagene). Selected plasmides were purified and sequenced as described earlier using T3, T7 and internal specific primers.

Northern blot analysis

Bovine chondrocyte mRNA was purified using a QuickPrep Micro mRNA Purification Kit (Pharmacia Biotech, Uppsala, Sweden), separated on a 1 % agarose-formaldehyde gel, transferred to nylon membranes and immobilized by UV crosslinking. cDNA-probes were ³²P-labelled with Random Primed DNA Labeling Kit (Boehringer Mannheim). Filters were prehybridized for 2-4 hours at 42 °C in 5x SSE, 5x Denharts solution, 0.1 % SDS, 50 µg/ml salmon sperm DNA and 50% formamide and then hybridized over night at 42 °C with the same solution containing the specific probe (0.5-1 x 10⁶ cpm/ml). Specifically bound cDNA-probes were analyzed using a phosphoimage system (Fuji). Filters were stripped by washing in 0.1% SDS, for 1h at 80°C prior to re-probing. The α 10-integrin cDNA-probe was isolated from the rac1-containing plasmid using the restriction enzymes BamHI (GIBCO-BRL) and NcoI (Boehringer Mannheim). The rat β 1-integrin cDNA probe was a kind gift from Staffan Johansson, Uppsala, Sweden (25).

Tissue staining

Human cartilage from the trochlear groove, obtained during surgery, was provided by Dr. Anders Lindahl, Sahlgrenska University Hospital, Gothenburg, Sweden. Frozen sections of cartilage tissue were fixed in acetone at -18°C for 5 minutes, washed in PBS and then treated

with 2mg/ml of hyaluronidase (Sigma) in PBS, pH 5.0 for 15 minutes at 37°C. After washing with PBS, sections were blocked for 15 minutes at room temperature in 1% H₂O₂ in PBS to remove endogenous peroxidase activity. Sections were then washed in PBS, blocked with 0.5% casein and 0.05% thimerosal in PBS (blocking buffer) for 15 minutes at room temperature and then incubated over night at 4°C with the affinity purified antibodies against the integrin subunits α 9 and α 10 (5 μ g/ml in blocking buffer). For control, α 10-antibody was preincubated with the α 10 peptide (0.1mg/ml) for 30 minutes at 4°C. After washing in PBS, sections were incubated with biotinylated goat anti-rabbit secondary antibody (Vector Laboratories Inc; diluted 1:200 in blocking buffer) at room temperature for 60 minutes. Washed sections were then incubated with vectastain ABC reagent (Vector Laboratories, Inc.) for 1 h at room temperature, washed, and the color was developed using 1 mg/ml of DAB in 0.1M Tris-HCl, pH 7.2 and 0.02 % H₂O₂. Sections were rinsed in water for 5 minutes followed by 75 % ethanol, 95 % ethanol and 99.5 % ethanol for 5 minutes each, and then three times in xylene for 3 minutes at room temperature. Samples were mounted in Pertex (HISTOLAB Products AB, Gothenburg, Sweden).

Results

Identification and isolation of the chondrocyte $\alpha 10$ integrin subunit

Affinity purification of ^{125}I -labelled membrane proteins from human chondrocytes on collagen type II-Sepharose followed by immunoprecipitation showed that these cells, in addition to $\alpha 1\beta 1$ and $\alpha 2\beta 1$, express an unidentified $\beta 1$ -related α subunit (Figure 1). This integrin subunit had a molecular weight of approximately 160 kD and was slightly larger than the $\alpha 2$ integrin subunit. This finding is in agreement with a previous study showing that bovine chondrocytes also express an unidentified collagen binding $\beta 1$ -associated α -subunit of similar M_r (25). To isolate this protein, we affinity purified collagen type II-binding proteins from bovine chondrocyte since. The cell lysate was applied to a fibronectin-Sepharose precolumn followed by a collagen type II-Sepharose column. As shown in Figure 2, a number of proteins were eluted from the affinity columns, however, a protein with M_r of approximately 160 kD was specifically eluted with EDTA from the collagen column but not from the fibronectin column. The M_r of this protein corresponded with the M_r of the unidentified $\beta 1$ -related integrin subunit (Figure 1). The 160 kD protein band was excised from the SDS-PAGE gel, digested with trypsin or Endoproteinase Lys C and several of the isolated peptides were analyzed. Table 1 shows the amino acid sequence of 6 individual peptides.

Cloning and sequencing of the human integrin α -subunit homologue

The nucleotide sequence of peptide 1 (Table 1) was obtained by PCR-amplification, cloning and sequencing of bovine cDNA. From this nucleotide sequence an exact primer was designed and then applied in PCR-amplification with degenerate primers corresponding to peptides 2-6 (Table I). Primers corresponding to peptides 1 and 2 amplified a 900 bp PCR-fragment from bovine cDNA which was cloned, sequenced and then used for screening of a human articular

chondrocyte λ ZapII cDNA library to obtain the human integrin α -subunit homologue. Two overlapping clones, hc1 and hc2 (Figure 3), were isolated, subcloned and sequenced. These clones contained 2/3 of the nucleotide sequence including the 3' end of the cDNA. A third clone (rac1; Figure 3), containing the 5' end of the α 10 cDNA, was obtained using the RACE technique. From these three overlapping clones of α 10 cDNA, 3884 nucleotides were sequenced (Figure 4). The sequence contains a 3504-nucleotide open reading frame that is predicted to encode a 1167 amino acid mature protein. The predicted sequence included a signal peptide (22 aa), a long extracellular domain (1098 aa), a transmembrane domain (25 aa) and a short cytoplasmic domain (22 aa). Sequence analysis of the 160 kD protein sequence showed that it was a member of the integrin α -subunit family and the protein was named α 10.

Comparison of α 10 integrin subunit with other α -subunits

Alignment of α 10 with known α subunits showed that its structure follows the conserved pattern of integrin α subunits (Figure 5). The extracellular domain contains a seven-fold repeated sequence including FG and GAP consensus sequences, three putative divalent cation-binding sites (DxD/NxD/NxxxD) and an I-domain of 199 amino acids. The protein contains 10 potential N-linked glycosylation sites (NXT/S). The calculated molecular weight is 153 kD if carbohydrate chains with an average molecular weight of 2.5 kD are assumed to attach to all 10 putative glycosylation sites. This is in agreement with the molecular weight of α 10 as judged by SDS-PAGE where the M_r was estimated to 160 kD. In contrast to most α -integrin-subunits the cytoplasmic domain does not contain the conserved sequence KxGFF (R/K) R (Table II). Instead the predicted amino acid sequence is KLGFFAH. The deduced amino acid sequence of α 10 showed the highest identity to the

collagen binding integrin subunits $\alpha 1$ (37%) and $\alpha 2$ (35%). The similarity of integrin α subunits are shown in Figure 6.

Expression of the $\alpha 10$ integrin subunit on chondrocytes

Northern blot analysis of mRNA from bovine chondrocytes showed that a human $\alpha 10$ cDNA-probe hybridized with a single mRNA of approximately 5.4 kb (Figure 7). As a comparison, a cDNA-probe corresponding to the integrin subunit $\beta 1$ was used. This cDNA-probe hybridized a mRNA-band of approximately 3.5 kb on the same filter. Translation of the $\alpha 10$ nucleotide sequence revealed an open reading frame of 3504 nucleotides (Figure 4) which indicates that around 2000 nucleotides in the mRNA is not translated.

To study expression of $\alpha 10$ on protein level, ^{125}I labelled membrane proteins from human chondrocytes were immunoprecipitated with polyclonal antibodies against the integrin subunits $\beta 1$, $\alpha 1$, $\alpha 2$, $\alpha 3$ and $\alpha 10$ (Figure 8). A polyclonal peptide antibody, raised against the cytoplasmic domain of $\alpha 10$, precipitated two protein bands with M_r of 160 kD and 125 kD under reducing conditions. The $\alpha 10$ associated β -chain migrated as the $\beta 1$ integrin subunit both under reducing and non-reducing conditions (Figures 8a,b).

Expression of $\alpha 10$ in cartilage was examined by immunostaining of human articular cartilage from the trochlear groove with the polyclonal $\alpha 10$ -antibody. As shown in Figure 9, this antibody specifically stained the chondrocytes in the cartilage tissue sections. The staining was completely abolished when the antibody was preincubated with the $\alpha 10$ -peptide. A control antibody against the $\alpha 9$ integrin subunit did not stain chondrocytes in the tissue sections (Figure 9).

Discussion

We show in this study that human chondrocytes express a novel, $\beta 1$ -associated, collagen type II-binding integrin, in addition to integrins $\alpha 1\beta 1$ and $\alpha 2\beta 1$. We have, in an earlier study, presented some evidence for that bovine chondrocytes and human chondrosarcoma cells also express an unknown integrin (25). Bovine chondrocytes were readily available and therefore used in the isolation of the unknown integrin subunit.

In the affinity purification experiment, fibronectin-Sepharose was used as a precolumn since earlier studies showed that fibronectin is not a ligand for this integrin (data not shown). As shown in Figure 2, several proteins were eluted from the affinity columns. It was difficult to interpret the protein pattern as typical integrin bands were not clearly distinguished on the SDS-PAGE gel. Based upon the findings that the unknown chondrocyte integrin α -subunit in the immunoprecipitation experiment (Figure 1) had a M_r of 160 kD, a protein with similar M_r , which was specifically eluted with EDTA from the collagen type II-column, was excised from the gel and used for peptide sequencing. The difficulty to distinguish integrin bands on the gel may be explained by that most integrins eluted from the columns were partially degraded although a mixture of protease inhibitors were included in the lysate buffer. The novel $\alpha 10$ integrin subunit appeared resistant to proteolytic degradation. The immunoprecipitation experiments showed that $\alpha 2$ and the new α integrin subunit have similar M_r s under reducing conditions (Figure 1). To avoid contamination of $\alpha 2$, the 160 kD-protein was excised from the SDS-PAGE gel as a very narrow band. It likely contained one protein since human homologues to all six bovine peptides (Table 1) that were isolated from the 160 kD protein were found in the predicted amino acid sequence of human $\alpha 10$ subunit (Figure 4).

The deduced amino acid sequence of $\alpha 10$ shares the general structure of the integrin α -subunits described in previously published reports (6-21). The large extracellular N-terminal

part of $\alpha 10$ contains a seven-fold repeated sequence which was recently predicted to fold into a β -propeller domain (32). The integrin subunit $\alpha 10$ contains three putative divalent cation-binding sites (DxD/NxD/NxxxD) (52), a single spanning transmembrane domain and a short cytoplasmic domain. In contrast to most α -integrin subunits the cytoplasmic domain of $\alpha 10$ does not contain the conserved sequence KxGFF (R/K) R. The predicted amino acid sequence in $\alpha 10$ is KLGFFAH. Several reports indicate that the integrin cytoplasmic domains are crucial in signal transduction (53) and that membrane-proximal regions of both α - and β -integrin cytoplasmic domains are involved in modulating conformation and affinity state of integrins (54-56). It is suggested that the GFFKR motif in α -chains are important for association of integrin subunits and for transport to the plasma membrane (57). The KxGFFKR domain has been shown to interact with the intracellular protein calreticulin (58) and interestingly, calreticulin-null embryonic stem cells are deficient in integrin-mediated cell adhesion (59). It is, in this context, tempting to speculate that the sequence KLGFFAH in $\alpha 10$ may have a key function in regulating the affinity between $\alpha 10\beta 1$ and collagen.

Integrin α subunits are known to share an overall identity of 20-40% (60). Sequence analysis showed that the $\alpha 10$ subunit is most closely related to the I-domain containing α -subunits (Figure 6) with the highest identity to $\alpha 1$ (37%) and $\alpha 2$ (35%). The integrins $\alpha 1\beta 1$ and $\alpha 2\beta 1$ are known receptors for both collagens and laminins (24;61;62) and we have also recently demonstrated that $\alpha 2\beta 1$ interacts with the cartilage protein chondroadherin (42). Since $\alpha 10\beta 1$ was isolated on a collagen type II-Sepharose we know that collagen type II is a ligand for $\alpha 10\beta 1$. We have also shown by affinity purification experiments that $\alpha 10\beta 1$ interacts with collagen type I (data not shown) but it remains to be seen whether laminin or chondroadherin are also ligands for this integrin.

We and others have shown previously that $\alpha 1\beta 1$ and $\alpha 2\beta 1$ are present on isolated chondrocytes although $\alpha 2\beta 1$ is found in only minor amounts (25). The peptide antibody that we raised against the cytoplasmic domain of $\alpha 10$ immunoprecipitated two proteins from human chondrocytes with M_r of 125 kD and 160 kD. The M_r of 160 kD correlates with the unidentified $\beta 1$ -associated α -subunit that was affinity purified on collagen type II-Sepharose. Immunohistochemistry using the $\alpha 10$ -antibody showed staining of the chondrocytes in tissue sections of human articular cartilage. The antibody staining was clearly specific since preincubation of the antibody with the $\alpha 10$ -peptide completely abolished the staining. An antibody against the integrin subunit $\alpha 9$ was used as a control in these experiments we (6). This integrin is a receptor for tenascin-C (63) and is not known to be present in cartilage. The immunohistochemical experiments also clearly demonstrate that $\alpha 10\beta 1$ is present not only on isolated cells but also in tissue sections of human cartilage. Taken together, we have isolated and characterized a novel collagen type II-binding integrin designated $\alpha 10\beta 1$. The $\alpha 10$ subunit was isolated from bovine chondrocytes and the human homologue was cloned and sequenced. Antibodies against the $\alpha 10$ -integrin subunit stained chondrocytes in tissue sections of articular cartilage indicating that $\alpha 10\beta 1$ is one of the collagen type II binding integrins expressed in cartilage. It is very likely that this integrin is important in the communication between chondrocytes and the extracellular matrix.

Acknowledgments

We are most grateful to Dick Heinegård, Anders Aspberg, Danny Tuckwell and Martin Humphries for valuable discussions and to Michael Bayliss for providing the human chondrocyte library.

Grants were obtained from the Swedish Medical Research Council, Anna-Greta Crafoord's Stiftelse, Crafoord's stiftelser, Gustav V's 80-års fond, Greta och Johan Kock's stiftelse, Kungliga fysiografiska sällskapets stiftelse, Riksföreningen mot reumatism, Magnus Bergvall's Stiftelse, Thelma Zoe'ga's fond and Alfred Österlund's Stiftelse.

References

1. Springer, T.A. (1990) *Nature* 346, 425-434
2. Ruoslahti, E. (1991) *J.Clin.Invest.* 87, 1-5
3. Hynes, R.O. (1992) *Cell* 69, 11-25
4. Hemler, M.E. (1988) *Immunol.Today* 9, 109-113
5. Yamada, K.M. (1991) *J.Biol.Chem.* 266, 12809-12812
6. Palmer, E.L., Ruegg, C., Ferrando, R., Pytela, R., and Sheppard, D. (1993) *J.Cell Biol.* 123, 1289-1297
7. Takada, Y., Elices, M.J., Crouse, C., and Hemler, M.E. (1989) *EMBO J.* 8, 1361-1368
8. Poncz, M., Eisman, R., Heidenreich, R., Silver, S.M., Vilaire, G., Surrey, S., Schwartz, E., and Bennett, J.S. (1987) *J.Biol.Chem.* 262, 8476-8482
9. Larson, R.S., Corbi, A.L., Berman, L., and Springer, T. (1989) *J.Cell Biol.* 108, 703-712
10. Corbi, A.L., Kishimoto, T.K., Miller, L.J., and Springer, T.A. (1988) *J.Biol.Chem.* 263, 12403-12411
11. Argraves, W.S., Suzuki, S., Arai, H., Thompson, K., Pierschbacher, M.D., and Ruoslahti, E. (1987) *J.Cell Biol.* 105, 1183-1190
12. Corbi, A.L., Miller, L.J., O'Connor, K., Larson, R.S., and Springer, T.A. (1987) *EMBO J.* 6, 4023-4028

13. Briesewitz, R., Epstein, M.R., and Marcantonio, E.E. (1993) *J.Biol.Chem.* **268**, 2989-2996

14. Ziober, B.L., Vu, M.P., Waleh, N., Crawford, J., Lin, C.S., and Kramer, R.H. (1993) *J.Biol.Chem.* **268**, 26773-26783
15. Hogervorst, F., Kuikman, I., van Kessel, A.G., and Sonnenberg, A. (1991) *Eur.J.Biochem.* **199**, 425-433
16. Takada, Y. and Hemler, M.E. (1989) *J.Cell Biol.* **109**, 397-407
17. Takada, Y., Murphy, E., Pil, P., Chen, C., Ginsberg, M.H., and Hemler, M.E. (1991) *J.Cell Biol.* **115**, 257-266
18. Van der Vieren, M., Le Trong, H., Wood, C.L., Moore, P.F., St.John, T., Staunton, D.E., and Gallatin, W.M. (1995) *Immunity.* **3**, 683-690
19. Schnapp, L.M., Breuss, J.M., Ramos, D.M., Sheppard, D., and Pytela, R. (1995) *J.Cell Sci.* **108**, 537-544
20. Shaw, S.K., Cepek, K.L., Murphy, E.A., Russell, G.J., Brenner, M.B., and Parker, C.M. (1994) *J.Biol.Chem.* **269**, 6016-6025
21. Suzuki, S., Argraves, W.S., Arai, H., Languino, L.R., Pierschbacher, M.D., and Ruoslahti, E. (1987) *J.Biol.Chem.* **262**, 14080-14085
22. Ignatius, M.J., Large, T.H., Houde, M., Tawil, J.W., Barton, A., Esch, F., Carbonetto, S., and Reichardt, L.F. (1990) *J.Cell Biol.* **111**, 709-720
23. Gullberg, D., Gehlsen, K.R., Turner, D.C., Åhlén, K., Zijenah, L.S., Barnes, M.J., and

Rubin, K. (1992) *EMBO J.* 11, 3865-3873

-
24. Staaaz, W.D., Rajpara, S.M., Wayner, E.A., Carter, W.G., and Santoro, S.A. (1989) *J.Cell Biol.* 108, 1917-1924
25. Holmvall, K., Camper, L., Johansson, S., Rubin, K., Kimura, J.H., and Lundgren-Åkerlund, E. (1995) *Exp.Cell Res.* 221, 496-503
26. Forsberg, E., Ek, B., Engström, Å., and Johansson, S. (1994) *Exp.Cell Res.* 213, 183-190
27. Wayner, E.A. and Carter, W.G. (1987) *J.Cell Biol.* 105, 1873-1884
28. Weitzman, J.B., Pasqualini, R., Takada, Y., and Hemler, M.E. (1993) *J.Biol.Chem.* 268, 8651-8657
29. Elices, M.J. and Hemler, M.E. (1989) *Proc.Natl.Acad.Sci.U.S.A.* 86, 9906-9910
30. Languino, L.R., Colella, S., Zanetti, A., Andrieux, A., Ryckewaert, J.J., Charon, M.H., Marchisio, P.C., Plow, E.F., Ginsberg, M.H., Marguerie, G., and et al (1989) *Blood* 73, 734-742
31. Tuckwell, D.S., Humphries, M.J., and Brass, A. (1994) *Cell Adhes.Comm.* 2, 385-402
32. Springer, T.A. (1997) *Proc.Natl.Acad.Sci.U.S.A.* 94, 65-72
33. Colombatti, A., Bonaldo, P., and Doliana, R. (1993) *Matrix* 13, 297-306
34. Lee, C.H., Bradley, G., and Ling, V. (1995) *Cell Growth Differ.* 6, 347-354
35. Calderwood, D.A., Tuckwell, D.S., and Humphries, M.J. (1995) *Biochem.Soc.Trans.*

36. Kern, A., Eble, J., Golbik, R., and Kuhn, K. (1993) *Eur.J.Biochem.* **215**, 151-159
37. Tuckwell, D.S., Reid, K.B., Barnes, M.J., and Humphries, M.J. (1996) *Eur.J.Biochem.* **241**, 732-739
38. Kamata, T. and Takada, Y. (1994) *J.Biol.Chem.* **269**, 26006-26010
39. Dürr, J., Goodman, S., Potocnik, A., von der Mark, H., and von der Mark, K. (1993) *Exp.Cell Res.* **207**, 235-244
40. Salter, D.M., Hughes, D.E., Simpson, R., and Gardner, D.L. (1992) *Br.J.Rheumatol.* **31**, 231-234
41. Woods, V.L.J., Schreck, P.J., Gesink, D.S., Pacheco, H.O., Amiel, D., Akeson, W.H., and Lotz, M. (1994) *Arthritis Rheum.* **37**, 537-544
42. Camper, L., Heinegård, D., and Lundgren-Åkerlund, E. (1997) *J.Cell Biol* **138**, 1159-1167
43. Hemler, M.E., Sanchez Madrid, F., Flotte, T.J., Krensky, A.M., Burakoff, S.J., Bhan, A.K., Springer, T.A., and Strominger, J.L. (1984) *J.Immunol.* **132**, 3011-3018
44. Bottger, B.A., Hedin, U., Johansson, S., and Thyberg, J. (1989) *Differentiation.* **41**, 158-167
45. Sommarin, Y. and Heinegård, D. (1983) *Biochem.J.* **214**, 777-784
46. Häuselmann, H.J., Aydelotte, M.B., Schumacher, B.L., Kuettner, K.E., Gitelis, S.H., and Thonar, E.J.M.A. (1992) *Matrix* **12**, 116-129

47. Miller, E.J. (1972) *Biochemistry* 11, 4903-4909
-
48. Wessel, D. and Flugge, U.I. (1984) *Anal.Biochem.* 138, 141-143
49. Blobel, G. and Dobberstein, B. (1975) *J.Cell Biol.* 67, 835-851
50. Hellman, U. (1997) in *Protein structure analysis. Preparation, characterization, and microsequencing* (Kamp, R.M., Choli-Papadopoulou, T., and Wittmann-Liebold, B., eds) pp. 97-104, Spriner-Verlag, Heidelberg
51. Charles, I.G., Palmer, R.M., Hickery, M.S., Bayliss, M.T., Chubb, A.P., Hall, V.S., Moss, D.W., and Moncada, S. (1993) *Proc.Natl.Acad.Sci.U.S.A.* 90, 11419-11423
52. Tuckwell, D.S., Brass, A., and Humphries, M.J. (1992) *Biochem.J.* 285, 325-331
53. Dedhar, S. and Hannigan, G.E. (1996) *Curr.Opin.Cell Biol.* 8, 657-669
54. Hughes, P.E., O'Toole, T.E., Ylanne, J., Shattil, S.J., and Ginsberg, M.H. (1995) *J.Biol.Chem.* 270, 12411-12417
55. Puzon McLaughlin, W., Yednock, T.A., and Takada, Y. (1996) *J.Biol.Chem.* 271, 16580-16585
56. O'Toole, T.E., Katagiri, Y., Faull, R.J., Peter, K., Tamura, R., Quaranta, V., Loftus, J.C., Shattil, S.J., and Ginsberg, M.H. (1994) *J.Cell Biol.* 124, 1047-1059
57. De Melker, A.A., Kramer, D., Kuikman, I., and Sonnenberg, A. (1997) *Biochem J* 529-537
58. Rojiani, M.V., Finlay, B.B., Gray, V., and Dedhar, S. (1991) *Biochemistry* 30, 9859-

-
59. Coppolino, M.G., Woodside, M.J., Demarex, N., Grinstein, S., St Arnaud, R., and Dedhar, S. (1997) *Nature* **386**, 843-847
60. Hynes, R.O. (1992) *Curr. Opin. Genet. Dev.* **2**, 621-624
61. Santoro, S.A. (1986) *Cell* **46**, 913-920
62. Languino, L.R., Gehlsen, K.R., Wayner, E., Carter, W.G., Engvall, E., and Ruoslahti, E. (1989) *J. Cell Biol.* **109**, 2455-2462
63. Yokosaki, Y., Monis, H., Chen, J., and Sheppard, D. (1996) *J. Biol. Chem.* **271**, 24144-24150

Figure Legends

Figure 1

Affinity purification and immunoprecipitation of collagen type II-binding integrins from triton X-100 lysate of ^{125}I -labeled human chondrocytes. The lanes show immunoprecipitation of integrins using monoclonal antibodies against the integrin subunits $\beta 1$ (P4C10), $\alpha 1$ (TS2/7), $\alpha 2$ (P1E6) and $\alpha 3$ (P1B5). The proteins eluted by EDTA from the collagen type II-Sepharose are shown in lane E. The proteins were separated by SDS-PAGE (4-12%) under nonreducing conditions and visualised using a phosphoimager.

Figure 2

Affinity purification of the $\alpha 10$ integrin subunit on collagen type II-Sepharose. A triton X-100 lysate of bovine chondrocytes (2.5×10^9 cells) was applied to a fibronectin-Sepharose precolumn followed by a collagen type II-Sepharose column. The lanes show EDTA-eluted proteins from the fibronectin-Sepharose (A), flow-through from the collagen type II-Sepharose column (B) and EDTA-eluted proteins from the collagen type II-Sepharose (C). The eluted proteins were precipitated by methanol/chloroform, separated by SDS-PAGE (4-12%) under reducing conditions and stained with coomassie blue. The 160 kD-protein with affinity for collagen type II is indicated with a arrow.

Figure 3

Schematic map of the sequencing strategy. The overlapping $\alpha 10$ clones hc1 and hc2 were obtained by screening a human articular chondrocyte library with a bovine $\alpha 10$ probe. The Race1 clone was obtained from human chondrocyte cDNA using the RACE technique. Arrows indicate the direction and extent of nucleotide sequencing.

Figure 4

Nucleotide sequence and deduced amino acid sequence of the human $\alpha 10$ integrin subunit. The

amino acid translation is under the first nucleotide of the corresponding codon. The signal peptide cleavage site is marked with a arrow, human homologues to bovine peptide sequences are underlined and the I-domain is boxed. Metal ion binding sites are indicated with a broken underline, potential N-glycosylation sites are indicated by an asterisk and the putative transmembrane domain is double underlined. The normally conserved cytoplasmic sequence is indicated by a dot and dashed broken underline. The sequence data is available from xxxxx under accession number xxxxxx.

Figure 5

Diagrammatic comparison of $\alpha 10$ with the general structure of integrin α -subunits. The conserved repeats 1-7 are indicated with boxes. An I-domain is found in some α -integrin subunits. Other α integrin subunits are posttranslationally cleaved near the transmembrane domain (TM). The integrin subunit $\alpha 10$ contains the 7 conserved repeats and an I domain located between repeat 2 and 3.

Figure 6

Sequence similarity between integrin α -subunits. The similarity tree was constructed by using the GCG software and the program "Pileup". Percent identities were calculated using the Jotun Hein algorithm provided in the Lasergene DNASTAR software. The similarity tree indicates three different subfamilies of α -integrin subunits; one subfamily that contains I-domains ($\alpha 1$, $\alpha 2$, αM , αX , αL and $\alpha 10$), one subfamily that is cleaved ($\alpha 3$, $\alpha 5$, $\alpha 6$, $\alpha 7$, $\alpha 8$, αv and αIIb) and one subfamily that neither contains I-domains nor is cleaved ($\alpha 4$, $\alpha 9$).

Figure 7

Northern blot analysis of $\alpha 10$ and $\beta 1$ mRNA. Bovine chondrocyte mRNA was hybridized with ^{32}P -labeled cDNA probes corresponding to the integrin subunits $\alpha 10$ or $\beta 1$. The $\alpha 10$ -probe hybridized to an mRNA of 5.4 kb and the $\beta 1$ -probe hybridized to an mRNA of 3.5 kb on the same filter.

Figure 8

~~Immunoprecipitation of the $\alpha 10$ integrin subunit from human chondrocytes. Triton X-100~~

lysates of ^{125}I -labelled human chondrocytes were immunoprecipitated with polyclonal antibodies against the integrin subunits $\beta 1$, $\alpha 1$, $\alpha 2$, $\alpha 3$ or $\alpha 10$. The immunoprecipitated proteins were separated by SDS-PAGE (4-12%) under reducing (a) and non-reducing conditions (b) and visualised using a phosphoimager.

Figure 9

Immunostaining of human articular cartilage. An antibody raised against the cytoplasmic domain of $\alpha 10$ (see Table II) stained the chondrocytes in tissue sections of human articular cartilage (A). The staining was depleted when the antibody was preincubated with the $\alpha 10$ -peptide (B). A control antibody recognising the $\alpha 9$ integrin subunit did not bind to the chondrocyte (C).

Table I

Amino acid sequences of peptides from bovine $\alpha 10$ -integrin. Peptides were isolated by in-gel digestion with trypsin and sequenced by Edman degradation.

Table II

Comparison of the cytoplasmic tails of I-domain containing integrin α -subunits. The underlined sequence in $\alpha 10$ represents the peptide that was used for antibody-production.

Peptide	Amino acid sequence
1	DNTAQTSAYIQYEPHISI
2	GPGHWDR
3	AAFDCSGQR
4	FAMGALPD
5	FTASLDEWTTAAR
6	VDASFRPQGXLAP

Q1 KIGFFKRPLKKKMEK

Q2 KLGFFKRKYEKMTKNFDEIDEETELSS

Q10 KIGFFAIEKKIPEEEKREIEKLEQ

Q1M KLGFFKRQYKIDMMSEGGPPGAEPQ

Q1X KVGFFKRQYKIDMMIEANGQIAPFNGTQTPSPSEKPKSEK

Q1L KVGFFKRNLKEKMEAGRGVPNGIPAEISQLASGQEAGDPGCLKPLIEKDSIESGGGKD

Q1E KCGFFKRKYQQINLESIRKAQIKSENILLEEN

Title: Integrin subunit $\alpha 10$

Inventor: Evy Lundgren-Åkelund, Department of Cell and Molecular Biology, Lund University, Sweden

Abstract


This invention relates to a novel integrin subunit $\alpha 10$ and the use of this integrin as a marker of chondrocytes, smooth muscle cells or endothelial cells. This invention also relates to the use of antibodies recognizing integrin subunit $\alpha 10$ or cDNA encoding for integrin $\alpha 10$ peptides in identifying integrin subunit $\alpha 10$ on chondrocytes, smooth muscle cells and endothelial cells and determining the differentiation state of the cells.

Background of the invention

See enclosed manuscript entiteled "Isolation, cloning and sequence analysis of a the collagen-binding integrin $\alpha 10\beta 1$ expressed on chondrocytes".

Claims

1. Integrin subunit $\alpha 10$ used as a marker of chondrocytes, smooth muscle cells or endothelial cells.
2. The use of integrin $\alpha 10$ antibodies or cDNA encoding for integrin $\alpha 10$ -peptides as markers of the integrin subunit $\alpha 10$ on chondrocytes, smooth muscle cells and endothelial cells.
3. A peptide consisting of the amino acid sequence of the cytoplasmic domain of integrin subunit $\alpha 10$, KLGFFAHKKIPEEEKREEKLEQ.
4. Polyclonal antibodies raised against the peptide claimed in 3.
5. The use of integrin $\alpha 10$ -antibodies or cDNA encoding integrin $\alpha 10$ peptides to determine differentiation-state of chondrocytes during development, in pathological conditions such as reumatoid arthritis and osteoarthritis, in tissue regeneration and in therapeutic and physiological repair of cartilage.
6. The use of integrin $\alpha 10$ -antibodies or cDNA encoding integrin $\alpha 10$ peptides to determine differentiation-state of smooth muscle cells in development and in pathological conditions such as atherosclerosis and inflammation.
7. The use of integrin $\alpha 10$ -antibodies or cDNA encoding integrin $\alpha 10$ -peptides to determine differentiation-state of endothelial cells during embryonic development, angiogenesis, or development of cancer.
8. Activation or blockage of the integrin subunit $\alpha 10$ to stimulate extracellular matrix syntesis and repair or to decrease extracellular matrix degradation.










Fig. 3



[illegible]








Fig. 8a,b

W 530-02

